

Scaling of the quasiparticle spectrum for d-wave superconductors

Steven H. Simon and Patrick A. Lee

Department of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139

(November 18, 1996)

In finite magnetic field H , the excitation spectrum of the low energy quasiparticles in a 2 dimensional d -wave superconductor exhibits a scaling with respect to $H^{1/2}$. This property can be used to calculate scaling relations for various physical quantities at low temperature T . As examples, we make predictions for the scaling behavior of the finite magnetic field specific heat, quasiparticle magnetic susceptibility, optical conductivity tensor, and thermal conductivity tensor. These predictions are compatible with existing experimental data. Most notably, the thermal Hall coefficient κ_{xy} measured by Krishana et al. in YBCO is found to scale as $\kappa_{xy} \sim T^2 F(\alpha T/H^{1/2})$ for $T \lesssim 30K$ in agreement with our predictions, where α is a constant and F is a scaling function.

PACS: 74.25.Fy, 74.72.-h, 74.25.Jb

Submitted To Physical Review Letters

The scientific community has been slowly coming to a consensus that the high T_c superconductors have a d -wave order parameter [1]. One of the major differences between these and conventional superconductors is that d -wave superconductors have gapless low energy excitations in certain directions in k -space, whereas s -wave superconductors are gapped. As a result, the low temperature behavior in the d -wave case can be quite different from that of conventional superconducting materials. Thus, in order to properly interpret experiments on these novel materials (and eventually develop a microscopic theory), we must have a clear understanding of the physics associated with a d -wave order parameter. In this work, we attempt to elucidate some of this physics by deriving scaling relations obeyed by the quasiparticle energies and eigenfunctions. Using these relations, we are then able to deduce scaling properties of a number of important physical quantities.

We begin our analysis with the Bogolubov [2] equations, $\mathcal{H}\psi = \epsilon\psi$ where $\psi = (u, v)$ is a Nambu 2-spinor whose components are the particle-like and hole-like part of the quasiparticle wavefunction respectively. Here,

$$\mathcal{H} = \begin{pmatrix} h + V - E_F & \hat{\Delta} \\ \hat{\Delta}^* & -h^* - V + E_F \end{pmatrix} \quad (1)$$

with E_F the Fermi energy, and h the kinetic part of the effective single particle Hamiltonian, and $V(\mathbf{r})$ is the effective disorder potential. As a model system we will choose to work with the effective one-particle Hamiltonian $h = \frac{(\mathbf{p}-\mathbf{A})^2}{2m}$ with m the electron effective mass. Although this neglects any direct Hartree or exchange pieces of the interaction, such pieces are thought to be relatively unimportant except in renormalizing m and V . (In this paper we have set the charge of the electron e , the speed of light c , and Planck's constant $\hbar$ all to unity.)

In Eq. 1, $\hat{\Delta}$ is the gap operator for spin singlet superconductivity defined as $\hat{\Delta}g(\mathbf{r}) = \int d\mathbf{r}' \Delta(\mathbf{r}, \mathbf{r}')g(\mathbf{r}')$ for any $g(\mathbf{r})$, where $\Delta(\mathbf{r}, \mathbf{r}') = -v(\mathbf{r} - \mathbf{r}')\langle\psi_\uparrow(\mathbf{r})\psi_\downarrow(\mathbf{r}')\rangle$ with v the inter-electron interaction. If we rewrite $\Delta(\mathbf{r}, \mathbf{r}')$ in terms of center of mass coordinate $\mathbf{R} = \frac{\mathbf{r}+\mathbf{r}'}{2}$ and relative coordinate $\mathbf{x} = \mathbf{r} - \mathbf{r}'$, then Fourier transform with

respect to $\mathbf{x}$, we can write the gap function as $\Delta(\mathbf{R}, \mathbf{k})$.

In this work we consider a two dimensional d -wave superconductor. It is believed that this accurately represents the high T_c materials. We choose to consider a gap function with pure d_{xy} symmetry rather than $d_{x^2-y^2}$ for notational simplicity. The final results for $d_{x^2-y^2}$ are identical. The gap function is written as $\Delta(\mathbf{R}, \mathbf{k}) = \Delta_{d_{xy}}(\mathbf{R})k_x k_y / (k_F)^2$. Shifting back to the coordinates $\mathbf{r}$ and $\mathbf{r}'$, then integrating by parts, the gap operator can be re-expressed as

$$\hat{\Delta} = \frac{1}{p_F^2} \{p_x, \{p_y, \Delta_{d_{xy}}(\mathbf{r})\}\} \quad (2)$$

where p_x and p_y are the components of the momentum operator, p_F is the Fermi momentum, and here the brackets represent the symmetrization, $\{a, b\} = \frac{1}{2}(ab + ba)$. The function $\Delta_{d_{xy}}$ is the d -wave order parameter used in Ginzburg-Landau theory [3]. We can then consider calculating $\Delta_{d_{xy}}$ in an inhomogeneous system by using a Ginzburg-Landau approach, then using $\Delta_{d_{xy}}$ in Eq. 1 to find the quasiparticle spectrum. We note that this approach is not fully self-consistent in the sense that we will not use the derived quasiparticle states to then recalculate the gap function.

For a homogeneous system there are gapless nodes on the Fermi surface at the points $\mathbf{p} = (\pm p_F, 0)$ and $\mathbf{p} = (0, \pm p_F)$ where $\hat{\Delta}$ vanishes. To study the low lying excitations near these points, we linearize the Hamiltonian. As an example, we consider linearizing around the point $\mathbf{p} = (p_F, 0)$. We write $\psi = e^{ik_F x} \tilde{\psi}$ such that we can recast the Bogolubov equations as $(\tilde{\mathcal{H}}_0 + \tilde{\mathcal{H}}_1)\tilde{\psi} = \epsilon\tilde{\psi}$ where $\tilde{\mathcal{H}}_0$ is the leading linearized term

$$\tilde{\mathcal{H}}_0 = \begin{pmatrix} v_F(p_x - A_x) + V & \frac{1}{p_F} \{p_y, \Delta_{d_{xy}}(\mathbf{r})\} \\ \frac{1}{p_F} \{p_y, \Delta_{d_{xy}}^*(\mathbf{r})\} & v_F(-p_x - A_x) - V \end{pmatrix} \quad (3)$$

and $\tilde{\mathcal{H}}_1$ is the remaining piece

$$\tilde{\mathcal{H}}_1 = \begin{pmatrix} h & \hat{\Delta} \\ \hat{\Delta}^* & -h^* \end{pmatrix}, \quad (4)$$

where $v_F = p_F/m$ is the Fermi velocity.

For sufficiently small energy excitations, $\tilde{\mathcal{H}}_0$ is much greater than $\tilde{\mathcal{H}}_1$ and it will be a reasonable approximation to neglect $\tilde{\mathcal{H}}_1$. To determine when this is a good approximation, we consider the homogeneous case of $\Delta_{d_{xy}}$ a real constant with $\mathbf{A} = 0$, and $V = 0$. We then find that $\tilde{\mathcal{H}}_0$ is just the Dirac Hamiltonian for massless fermions in two dimensions, and thus has a conical linear spectrum of quasiparticles $\epsilon_{\mathbf{p}} = \pm \mathcal{E}(\mathbf{p})$ with $\mathcal{E}(\mathbf{p}) = \sqrt{(v_F p_x)^2 + (\Delta_{d_{xy}} p_y / p_F)^2}$. Note that the conical spectrum is highly anisotropic since $v_F \gg \Delta_{d_{xy}} / p_F$. For excitations at temperature T , the typical momenta are $p_x \sim T / v_F$ and $p_y \sim T p_F / \Delta_{d_{xy}}$. The largest term in $\tilde{\mathcal{H}}_1$ is then the term $p_y^2 / (2m)$ which would be on order $E_F (T / \Delta_{d_{xy}})^2$. For YBCO, photoemission spectroscopy [5] indicates that $E_F \approx 3,000K$, and $\Delta_{d_{xy}} \approx 450K$. In YBCO, we also note that the Fermi surface is not circular, but is somewhat flattened at the nodes (more square-like with rounded corners). This means we should really use an effective mass m_y in the $p_y^2 / (2m)$ term of $\tilde{\mathcal{H}}_1$ which lowers the energy scale of $\tilde{\mathcal{H}}_1$ by another factor of perhaps two or three. Thus, we estimate that $\tilde{\mathcal{H}}_1 / \tilde{\mathcal{H}}_0 \approx T / (200K)$, so that the condition $\tilde{\mathcal{H}}_0 \gg \tilde{\mathcal{H}}_1$ may be well satisfied at temperatures as high as $30K$.

We note that in Eq. 3 a nonzero A_x acts as a scalar potential for the quasiparticles. If we consider the case of a uniform superfluid velocity, we can choose a gauge where $\Delta_{d_{xy}}$ is a real constant and $\mathbf{A}$ is proportional to the velocity (i.e., London gauge). In this case, A_x in Eq. 3 acts as a scalar potential to yield a Doppler shifted spectrum $\epsilon_{\mathbf{p}} = v_F A_x \pm \mathcal{E}(\mathbf{p})$. Of course, were we to consider quasiparticles near the opposite $(-p_F, 0)$ node, we would have a spectrum $\epsilon_{\mathbf{p}} = -v_F A_x \pm \mathcal{E}(\mathbf{p})$. As pointed out by Volovik [6], in a magnetic field, screening currents exist with a typical velocity proportional to $H^{1/2}$ so that in a semiclassical approximation, the density of states at zero energy is proportional to $H^{1/2}$.

Let us now apply a magnetic field H perpendicular to the plane of the sample to create a vortex lattice such that the phase of the gap $\Delta_{d_{xy}}$ twists a full 2π as we go around each vortex. (Since the screening length is very long, we can assume H is homogeneous). The distance between the vortices is proportional to the magnetic length $l_H \sim H^{-1/2}$. We now claim that at low T , to a very good approximation, the Hamiltonian $\tilde{\mathcal{H}}_0$ has a simple scaling form that we write (in a slight abuse of notation) as $\tilde{\mathcal{H}}_0^H(\mathbf{r}) = [H/H_0]^{1/2} \tilde{\mathcal{H}}_0^{H_0}(\mathbf{r}[H/H_0]^{1/2})$. In other words, if we can find the eigenvectors $\tilde{\psi}_n^{H_0}(\mathbf{r})$ and eigenenergies $\epsilon_n^{H_0}$ of the Hamiltonian $\tilde{\mathcal{H}}_0$ in field H_0 , then the eigenenergies and eigenvectors in field H can be written as

$$\tilde{\psi}_n^H(\mathbf{r}) = \tilde{\psi}_n^{H_0}(\mathbf{r}[H/H_0]^{1/2}) \quad (5)$$

$$\epsilon_n^H = [H/H_0]^{1/2} \epsilon_n^{H_0}. \quad (6)$$

The first of these equations is the statement that the functional form of the eigenvector scales as the vortex lattice, whereas the second is just a reflection of the Hamil-

tonian being linear in momentum. In order to demonstrate that these scaling properties hold, we consider each term in $\tilde{\mathcal{H}}_0$ individually. It easy to show that the vector potential in a field H can be written in a scaling form $\mathbf{A}_H(\mathbf{r}) = [H/H_0]^{1/2} \mathbf{A}_{H_0}(\mathbf{r}[H/H_0]^{1/2})$, and similarly, $\mathbf{p}$ must scale as the inverse of the characteristic length l_H so that $\mathbf{p}^H = [H/H_0]^{1/2} \mathbf{p}^{H_0}$. Thus we need only examine V and $\Delta_{d_{xy}}$.

We first consider the scaling of $\Delta_{d_{xy}}$. In order to have the desired scaling of $\tilde{\mathcal{H}}_0$, we must have $\Delta_{d_{xy}}^H(\mathbf{r}) = \Delta_{d_{xy}}^{H_0}(\mathbf{r}[H/H_0]^{1/2})$. This is simply the statement that, like the wavefunction $\tilde{\psi}$, the functional dependence of $\Delta_{d_{xy}}$ on position scales with the vortex lattice. This assumption is clearly not correct near the vortex cores where there is some fixed width to the core that is relatively independent of field. However, away from the cores, for $T \ll T_c$, the magnitude of $\Delta_{d_{xy}}$ is fixed and only the phase varies. For $|\Delta_{d_{xy}}|$ fixed, it can be shown that the function form of the phase that minimizes the Ginzburg-Landau free energy has the correct scaling properties. Thus, for $T \ll T_c$, we expect that Hamiltonian will scale to a very good approximation.

We note that in general, currents in d -wave superconductors can introduce some small amount of s -wave component (Δ_s) to the order parameter [4]. It can also be shown that the above scaling laws hold to linear order in $\Delta_s / \Delta_{d_{xy}}$, so that for $\Delta_{d_{xy}} \gg \Delta_s$ the scaling laws (Eqs. 5 and 6) are preserved [7].

Finally, we turn to consider the disorder term V . For Gaussian delta-function correlated disorder such that $\langle V \rangle = 0$, and $\langle V(\mathbf{r})V(\mathbf{r}') \rangle = V_0 \delta(\mathbf{r} - \mathbf{r}')$, the disorder does not define a length scale so that given a realization of disorder $V(\mathbf{r})$, another configuration $V'(\mathbf{r}) = [H/H_0]^{1/2} V(\mathbf{r}[H/H_0]^{1/2})$ is equally likely. In other words, the disorder term (in an ensemble average) has the proper scaling properties to preserve Eqns. 5 and 6. For more general disorder with a nonzero correlation length, these scaling laws will no longer hold precisely. We note, however, that since there is no Anderson's theorem [2] for d -wave superconductors, the introduction of disorder will reduce the overall value of the gap and thereby reduce the maximum temperature at which the condition $\tilde{\mathcal{H}}_0 \gg \tilde{\mathcal{H}}_1$ is satisfied.

Neglecting disorder once again, $\Delta_{d_{xy}}$ and $\mathbf{A}$ can both be considered to be periodic functions with the periodicity of the vortex lattice [7]. Due to this periodicity, the eigenstates can be divided into Brillouin zones with one band of excitations per zone. The first zone should have a maximum k -vector of approximately $|\mathbf{k}_{\max}| \approx l_H^{-1}$, with l_H the magnetic length. The number of different zones with momentum less than some k is roughly $(k/k_{\max})^2$. The typical energy scale of an excitation of wavevector k is $k \sqrt{\Delta_{d_{xy}} v_F / p_F}$. Thus, the typical energy E_n of the n^{th} band is given roughly by $E_n \sim k_{\max} \sqrt{n \Delta_{d_{xy}} / m} \sim \sqrt{n \omega_c \Delta_{d_{xy}}}$ where $\omega_c = B/m$ is the cyclotron frequency. It is then convenient to define the dimensionful con-

stant $\alpha = \sqrt{m/\Delta_{xy}}$ such that $E_n^2 \approx n\alpha^{-2}B$. Finally, it will be useful to define the dimensionless parameter $x = \alpha T/H^{1/2}$ which is roughly the number squared of bands that are considerably occupied at temperature T . For YBCO, $\alpha \approx .05 \text{ Tesla}^{1/2}/K$.

Using the above described scaling laws we can extract a number of important statements about physical quantities. As a first example we examine the specific heat. We write the energy as

$$U = \sum_n \epsilon_n^H f(\epsilon_n^H/T) \\ = [H/H_0]^{1/2} \sum_n \epsilon_n^{H_0} f(\epsilon_n^{H_0}[H/H_0]^{1/2}/T) \quad (7)$$

where f is the Fermi function. The volume ν of the system here scales as $l_H^2 \sim 1/H$ so that $\nu = \nu_0[H_0/H]$. Thus, the energy density can be written as $U/\nu = H^{3/2}F_U(\alpha T/H^{1/2})$ where F_U is some scaling function that we can write down in terms of eigenenergies, but can not completely evaluate without fully diagonalizing $\tilde{\mathcal{H}}_0$. Here we have used the fact that the sum in Eq. 7 is only a function of the dimensionless quantity $x = \alpha T/H^{1/2}$. Note that here and elsewhere in this paper, we will assume that we are sufficiently far below T_c that the magnitude of the gap does not change much with T . Differentiating to obtain the electronic specific heat per unit volume we find that

$$C_v = TH^{1/2}F_C(\alpha T/H^{1/2}) \quad (8)$$

where F_C is again some unknown scaling function. We note that this scaling form does not include contributions to the specific heat from normal state electrons in the vortex cores. These contributions, however, are thought to be small [6]. As discussed above, in a semiclassical approximation, the density of states at zero energy is proportional to $H^{1/2}$ so that the specific heat is proportional to $TH^{1/2}$ at low temperatures as predicted by Volovik [6]. Thus the function F_C should be a constant at small argument. Deviations from this constant should first occur when we start to fill more than one band. This happens at $x \approx 1$, or at $(T/H^{1/2}) \approx 20 \text{ K/Tesla}^{1/2}$.

Similarly, starting with the free energy $F = U - TS$, where $S = \sum_n (f_n \ln f_n + (1 - f_n) \ln(1 - f_n))$ and $f_n = f(\epsilon_n/T)$ we can write the free energy per unit volume in the scaling form $F/\nu = H^{3/2}F_F(T/H^{1/2})$. We then conclude that the quasiparticle magnetic susceptibility per unit volume scales as $\chi = d^2(F/\nu)/dM^2 = \frac{1}{T}F_\chi(\alpha T/H^{1/2})$ with F_χ an unknown scaling function. Since there is no crossing of states through the Fermi level as we change magnetic field, we do not predict any de Haas-van Alphen oscillations. However, there may be oscillatory contributions to the susceptibility from the condensed fraction and the normal vortex cores that we do not consider here [9].

We now turn to consider electrical and thermal transport properties. We first define [11] the charge velocity

operator $\mathbf{v}^{(1)} \equiv i[\mathcal{H}, \mathbf{r}\sigma_z]$ and the thermal velocity operator $\mathbf{v}^{(2)} \equiv i[\mathcal{H}, \{\mathcal{H}, \mathbf{r}\}]$. Operating on a state $\psi = e^{ik_F x} \tilde{\psi}$ near the node at $(p_F, 0)$, we find

$$\mathbf{v}^{(1)} e^{ik_F x} \tilde{\psi} = e^{ik_F x} [\tilde{\mathcal{H}}_0, \{\sigma_z, \mathbf{r}\}] \tilde{\psi} + \text{smaller terms} \quad (9)$$

$$\mathbf{v}^{(2)} e^{ik_F x} \tilde{\psi} = e^{ik_F x} [\tilde{\mathcal{H}}_0, \{\tilde{\mathcal{H}}_0, \mathbf{r}\}] \tilde{\psi} + \text{smaller terms} \quad (10)$$

where the smaller terms are typically smaller by order $\tilde{\mathcal{H}}_1/\tilde{\mathcal{H}}_0$. It is then easy to see from this form that the operator $\mathbf{v}^{(2)}$ scales as $H^{1/2}$ whereas the operator $\mathbf{v}^{(1)}$ scales as H^0 .

We now use the Kubo formula to write the generalized response function at frequency ω by [11]

$$L_{ij}^{ab} = \frac{T}{\nu} \sum_{nm} \frac{\langle n | \mathbf{v}_i^{(a)} | m \rangle \langle m | \mathbf{v}_j^{(b)} | n \rangle f(\epsilon_n/T, \epsilon_m/T)}{(\epsilon_n - \epsilon_m - \omega - i0^+)(\epsilon_n - \epsilon_m + i0^+)}$$

where f is the thermal occupation factor, ν is the volume of the system, the indices i and j take the values $\hat{\mathbf{x}}$ and $\hat{\mathbf{y}}$, and the indices a and b take the values 1 and 2 (for charge and heat transport respectively). Noting that the volume of the system scales as H^{-1} and the energies all scale as $H^{1/2}$ we immediately obtain the two parameter scaling law $L_{ij}^{ab} \sim T^{a+b-1} F_{ij}^{ab}(\alpha T/H^{1/2}, \alpha \omega/H^{1/2})$, where F_{ij}^{ab} is again some scaling function that we will not evaluate. The real part of the optical conductivity tensor is defined as $\text{Re}[\sigma_{ij}] = \frac{1}{T} \text{Re}[L_{ij}^{11}]$ which immediately yields a two parameter scaling law for the optical conductivity.

$$\text{Re}[\sigma_{ij}] \sim F_{ij}^{11}(\alpha T/H^{1/2}, \alpha \omega/H^{1/2}). \quad (11)$$

It should be noted that in this Kubo formula calculation the response of the superfluid fraction has been neglected. This then does not include, for example, the response of the system due to the motion of vortices [12].

We now turn our attention to the DC ($\omega = 0$) thermal conductivity tensor κ , defined as the matrix that relates the heat current $\mathbf{j}_q$ to the temperature gradient via $\mathbf{j}_q = \kappa \nabla T$. We note that experimentally, a large part of the diagonal components of this tensor are due to phonon transport of heat. However, the Hall (off diagonal) component of this tensor should be completely electronic in origin [10]. Note, that when calculating κ , one must usually take into account the effect of the thermoelectric coefficient L^{12} . However, here we can neglect that contribution, since there is never any voltage in the superconducting state. Thus, we have $\kappa_{ij} = \frac{1}{T^2} L_{ij}^{22}$, and we obtain the naive scaling law $\kappa_{ij} \sim T F_{ij}^{22}(\alpha T/H^{1/2})$. Although this is indeed the correct scaling form for the (electronic part of the) diagonal component of the tensor, it is not correct for the Hall component. It can in fact be shown that the scaling function F_{xy}^{22} here is precisely zero due to the particle-hole symmetry inherent in the linearized Hamiltonian $\tilde{\mathcal{H}}_0$. This result is very easy to understand. Imposing a heat source on one side of the system excites many particles and holes. Both particles and holes diffuse in the direction of the heat sink. In a




FIG. 1. Thermal Hall data from reference [13]. (a) Thermal Hall transport coefficient κ_{xy} plotted against external magnetic field H at temperatures (from bottom to top) 20K, 22.5K, 25K, 27.5K, and 30K. For technical reasons data has not yet been taken below 20K. (b) Collapse of these five curves according to the scaling law shown in Eq. 12. Note that the characteristic scale of $H^{1/2}/T$ is approximately .05 Tesla^{1/2}/K.

magnetic field, the particles curve one way and the holes curve the other way. Thus, when there is particle-hole symmetry, there is no net Hall transport of heat.

The proof that the linearized Hamiltonian yields $\kappa_{xy} = 0$ is a little bit involved. The particle-hole symmetry is expressed mathematically by saying that given an eigenpair $\epsilon_n, \tilde{\psi}_n$ satisfying $\tilde{\mathcal{H}}_0 \tilde{\psi}_n = \epsilon_n \tilde{\psi}_n$, it can be shown that there exists another eigenvector $\tilde{\psi}'_n = \sigma_y \tilde{\psi}_n^*$ with the same eigenvalue also satisfying $\tilde{\mathcal{H}}_0 \tilde{\psi}'_n = \epsilon_n \tilde{\psi}'_n$ where σ_y is the usual Pauli spin matrix. Using this symmetry and adding up the contributions to κ_{xy} from all four nodes on the Fermi surface, it can be shown that κ_{xy} vanishes. It should be noted that κ_{xy} remains zero even when we include the smaller terms in from Eq. 10. We also note that this mechanism does not force $\sigma_{xy} = 0$. Further details of this calculation will be published elsewhere [7].

In order to find a nonzero κ_{xy} , we must break the particle-hole symmetry by including the contributions from $\tilde{\mathcal{H}}_1$. As mentioned above, at low enough T , we have $\tilde{\mathcal{H}}_1 \ll \tilde{\mathcal{H}}_0$ so that this term can be treated perturbatively. Inclusion of this term then shifts the eigenenergies via $\epsilon_n \rightarrow \epsilon_n + \delta\epsilon_n$ and the eigenstates $|n\rangle \rightarrow |n\rangle + \delta|n\rangle$. To lowest order these shifts are given by the usual expressions from perturbation theory which then obtain the following scaling forms

$$\delta\epsilon_n^H = \langle n^H | \tilde{\mathcal{H}}_1 | n^H \rangle = [H/H_0]^{1/2} \delta\epsilon_n^{H_0}$$

$$\delta|n^H\rangle = \sum_m \frac{|m^H\rangle \langle m^H | \tilde{\mathcal{H}}_1 | n^H \rangle}{\epsilon_n^H - \epsilon_m^H} = [H/H_0]^{1/2} \delta|n^{H_0}\rangle.$$

Note that both of these corrections scale as $H^{1/2}$. Including these first order corrections into the Kubo formula and expanding we find these correction terms give a leading contribution to the thermal Hall conductivity that scales as

$$\kappa_{xy} \sim T^2 F_{\kappa_{xy}}(\alpha T/H^{1/2}) \quad (12)$$

with $F_{\kappa_{xy}}$ again some scaling function. As shown in Fig. 1, experimental results of reference [13] do indeed show this scaling form at temperatures below 30K (For technical reasons data has not yet been taken at temperatures below 20K). As discussed above, the characteristic scale for features in the function $F_{\kappa_{xy}}$ (i.e., where the curve becomes nonlinear) should be seen at $x \approx 1$ or $H^{1/2}/T \approx .05$ Tesla^{1/2}/K, which is in good agreement with experiment.

In conclusion, we have found that the scaling properties of the quasiparticle spectrum in d -wave superconductors provides a very general and powerful tool for analyzing various physical quantities.

It is a pleasure to acknowledge very helpful conversations with K. Matveev, D. K. K. Lee, and particularly L. S. Levitov. The authors are greatly indebted to N. P. Ong and K. Krishana for sending us their experimental data prior to publication. This work was supported by NSF Grant Number DMR-9523361.

-
- [1] See D. J. Van Harlingen, Rev. Mod. Phys. **67**, 515 (1995); C. C. Tsuei et al., Phys. Rev. Lett. **73**, 593 (1994).
 - [2] See, for example, P. G. de Gennes, *Superconductivity of Metals and Alloys*, (Addison-Wesley; Reading, Mass, 1966).
 - [3] D. L. Feder and C. Kallin, preprint cond-mat/9609248; J. Xu, Y. Ren, and C. S. Ting, Phys. Rev. B, **52**, 7663 (1995).
 - [4] See for example, M. Franz, et al., Phys Rev B **53**, 5795 (1996) and references therein.
 - [5] Z. X. Shen and D. S. Dessau, Phys. Rep. **253**, 1 (1995).
 - [6] G. E. Volovik, JETP Letters, **58**, 470 (1993).
 - [7] S. H. Simon, in preparation.
 - [8] K. A. Moler, et al., Phys. Rev. Lett. **73**, 2744 (1994); J. Phys. Chem. Solids **56**, 1899 (1995); R. A. Fisher et al., Physica C, **252**, 237 (1996).
 - [9] See for example M. R. Norman, A. H. MacDonald, and H. Akera, Phys. Rev. B **51**, 5927 (1995).
 - [10] K. Krishana, J. M. Harris, and N. P. Ong, Phys. Rev. Lett. **75**, 3529 (1995).
 - [11] G. D. Mahan, *Many-Particle Physics*, 2ed., (Plenum, NY 1990).
 - [12] H. D. Drew, et al., preprint; H.-T.S. Lihn, et al., Phys. Rev. Lett. **76**, 3810 (1996).
 - [13] K. Krishana and N. P. Ong, unpublished.